



HAL
open science

Cerebral oxygenation and cognitive performance under acute normobaric hypoxia

Eléonore Fresnel, Gérard Dray, Simon Pla, Guilhem Belda, S. Perrey

► **To cite this version:**

Eléonore Fresnel, Gérard Dray, Simon Pla, Guilhem Belda, S. Perrey. Cerebral oxygenation and cognitive performance under acute normobaric hypoxia. ACAPS 2021 - 19ème congrès international des chercheurs en Activités Physiques et Sportives, Oct 2021, Montpellier, France. ACAPS 2021 - 19ème congrès international des chercheurs en Activités Physiques et Sportives, 2021. hal-03443395

HAL Id: hal-03443395

<https://hal.mines-ales.fr/hal-03443395>

Submitted on 2 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cerebral oxygenation and cognitive performance under acute normobaric hypoxia

01 INTRODUCTION

► **Altitude hypoxia** can **degrade executive functions and associated behavioral responses** in fighter pilots and so be at the origin of accidents in flight.

► **Working Memory (WM)** is impaired during exposure to hypoxia beyond 3000m (Malle et al., 2013)

► **Cerebral oxygenation responses** are correlated to cognitive performance during hypoxia (Williams et al., 2019)

Purpose

Identify the effects of acute exposure in normobaric hypoxia on cerebral oxygenation and cognitive performance during a working memory task

02 METHODS

Normoxia | Hypoxia
FiO₂ = 21 % | FiO₂ = 11.3%




Fig.1: Experimental set-up

Participants:

- 10 healthy adults attended 2 randomized sessions:
- 1 normoxia (N)
- 1 hypoxia (H)

Design (Fig.1):

- 9 blocks of 49 stimuli of a WM task (2-back)
- N session: 9 blocks under N
- H session: 2 blocks N (T1) → 3 blocks up to H (T2) → 4 blocks H (T3)

Variables:

- Cerebral oxygenation (PortaLite, Artinis; 10Hz): oxygenated (O₂Hb) and desoxygenated (HHb) blood
- Working memory performance and reaction time (RT)
- Nasa-TLX (subjective questionnaire)

RESULTS 03




Fig.2: Performance of Working Memory task depending on condition (H vs N) and time periods




Fig.3: Cerebral oxygenation under N and H during the cognitive task

IN BRIEF

Working Memory (Fig.2)

- No difference of condition (H / N) on performance and RT
- No difference of time on performance and RT

Cerebral oxygenation (Fig.3)

- Difference between N and H on O₂Hb and HHb
- ↘ O₂Hb and ↗ HHb with time in H

NASA-TLX

- Hypoxia ↗ perceptive workload (p < .05)

04 CONCLUSION

► Hypoxia (~5000m) induced reduction in cerebral oxygenation but not with cognitive performance.

The difficulty of WM might be too low in altering the cognitive performance.

► Other executive functions must be taken in consideration (e.g., inhibition and selective attention with the Stroop task)

Notes

¹ EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Alès, Montpellier, France

² Semaxone, Rochefort du Gard (30650), France

References

- Malle C, et al. (2013). Aviat Space Environ Med. DOI: 10.3357/ASEM.3482.2013
- Williams TB, et al. (2019). Exp Physio. DOI: 10.1113/EP087647

Acknowledgements

This work was publicly funded through ANR (the French National Research Agency) under the "Investissements d'avenir" program with the reference ANR-16-IDEX-0006.

Contact

eleonore.fresnel@gmail.com
stephane.perrey@umontpellier.fr