



HAL
open science

The reserve of joint torque determines movement coordination

Germain Faity, Denis Mottet, Simon Pla, Jérôme Froger

► **To cite this version:**

Germain Faity, Denis Mottet, Simon Pla, Jérôme Froger. The reserve of joint torque determines movement coordination. *Scientific Reports*, 2021, 11 (1), pp.23008. 10.1038/s41598-021-02338-4 . hal-03583855

HAL Id: hal-03583855

<https://imt-mines-ales.hal.science/hal-03583855>

Submitted on 22 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OPEN

The reserve of joint torque determines movement coordination

Germain Faity¹, Denis Mottet^{1✉}, Simon Pla¹ & Jérôme Froger²

Humans coordinate biomechanical degrees of freedom to perform tasks at minimum cost. When reaching a target from a seated position, the trunk-arm-forearm coordination moves the hand to the well-defined spatial goal, while typically minimising hand jerk and trunk motion. However, due to fatigue or stroke, people visibly move the trunk more, and it is unclear what cost can account for this. Here we show that people recruit their trunk when the torque at the shoulder is too close to the maximum. We asked 26 healthy participants to reach a target while seated and we found that the trunk contribution to hand displacement increases from 11 to 27% when an additional load is handled. By flexing and rotating the trunk, participants spontaneously increase the reserve of anti-gravitational torque at the shoulder from 25 to 40% of maximal voluntary torque. Our findings provide hints on how to include the reserve of torque in the cost function of optimal control models of human coordination in healthy fatigued persons or in stroke victims.

When reaching to grasp an object from a seated position, the body segments that bring the hand closer to the target are the trunk, arm and forearm. This kinematic chain is redundant so that, theoretically, reaching the final posture can be achieved in an endless number of different patterns of coordination of the trunk, arm and forearm¹. However, empirically, the reaching movement quasi exclusively relies on shoulder flexion and elbow extension².

According to the optimal control framework, the CNS achieves goals at a minimal cost³. To coordinate body segments, the CNS minimises both error and effort⁴, where the error is task related and the effort includes internal costs^{5,6}, such as minimum energy expenditure, minimum muscle fatigue or minimum sense of effort⁷. When reaching to grasp an object from a seated position, the CNS takes into account that the trunk is massive, with greater inertia than the upper arm and forearm, and with a larger lever arm that requires more torque to start and stop. Thus, when it can do otherwise, the CNS has no interest in recruiting the costly trunk within the coordination.

However, the CNS sometimes recruits the trunk. For example, the trunk is recruited when the target is too far away to be reached with the shoulder-elbow coordination⁸. Obviously, to minimise the error, one has to pay the price of a higher effort. More surprisingly, the trunk is also recruited when the target is not far away, typically in people suffering from hemiparesis following a stroke^{9,10}, in children with and without cerebral palsy¹¹ or in fatigued healthy people¹². Peeters et al. (2018) argue that it may be easier to accurately perform a reaching task in children or patients with cerebral palsy when the arm is not fully extended due to a smaller lever arm at the shoulder, which makes sense because forward trunk flexion decreases the antigravitational torque at the shoulder in the final posture¹³. Fuller et al. (2013) suggest a similar hypothesis for healthy subjects who begin to use trunk flexion after a fatigue protocol. Using the trunk would relieve the shoulder muscles by recruiting the trunk muscles. Hence, it is likely that the trunk is recruited to limit the torque at the shoulder joint in the final posture. Doing so, the objective of the task would be pursued (i.e. bringing the hand closer to the target), while the effort would be better distributed and with greater torque reserve for the involved joints.

In this paper, we hypothesise that trunk recruitment becomes significant when the shoulder is too close to maximal voluntary torque. We test this hypothesis by comparing the final posture when the hand is unloaded versus when the hand is loaded at 75% of the maximum antigravity shoulder torque.

¹EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Alès, Montpellier, France. ²EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Alès, CHU Nîmes, Le Grau du Roi, France. ✉email: denis.mottet@umontpellier.fr

Methods

Participants. Twenty-six healthy participants (12 males, age 21 ± 3 years, 3 left-handed, height 1.73 ± 0.09 m, weight 66.92 ± 9.29 kg, maximum voluntary force (MVF) of shoulder flexion 65.32 ± 19.50 N m) participated in this study. Participants were excluded if they had shoulder pain or any other problem that could affect the reaching task. Written informed consent was obtained from all participants before their inclusion. This study was performed in accordance with the 1964 Declaration of Helsinki. The Institutional Review Board of EuroMov at the University of Montpellier approved the study (IRB-EM 1901C).

Procedure. Participants had to reach a target with the side of their thumb nail. The target was table tennis ball fixed in front of the participant at a height of 80 cm¹⁴. The distance to the target was adjusted to correspond to the length of the participant's active stretched arm measured from the medial axilla to the distal wrist crease¹⁵. The height of the chair was adjusted so that the target horizontal projection was between the participant's navel and xiphoid process.

In the starting position, the participants were seated with their feet on the floor, their backs in contact with the back of the chair and their forearms on the armrest. The reaching movement was assessed in 2×2 conditions: spontaneous trunk use or restrained trunk use \times hand unloaded or hand loaded. In the spontaneous trunk use condition, participants had to reach the target at a natural pace, wait 1 s and return to the starting position. In the restrained trunk use condition, participants had to reach the target while minimising trunk movement: the experimenter manually applied a light proprioceptive feedback on the participant's shoulders, as a reminder to minimise trunk movement. We did not use a belt to restrain the trunk in order to leave the participant free to use the trunk if necessary, and thus avoid task failure. To avoid a loss of spontaneity, the spontaneous condition was always performed before the restraint condition¹⁴. Participants were given a couple of minutes of rest between conditions to avoid a fatigue effect.

In both trunk conditions, participants performed 5 unloaded trials and 5 loaded trials, i.e., carrying a dumbbell (Fig. 1). The assessed hand was chosen pseudo-randomly (12 left, 14 right) so that half of the participants performed the task with their dominant hand, and the other half with their non-dominant hand. In the hand unloaded condition, the weight of the arm without the dumbbell corresponded to 13.0% ($\pm 3.7\%$) of the MVF of the shoulder. In the hand loaded condition, the weight of the dumbbell was set as a function of the MVF of the participant, in order that the weight of the arm including the dumbbell corresponded to 75.0% ($\pm 5.5\%$) of the MVF of the shoulder.

The MVF of the shoulder was the maximum torque producible by the shoulder during an isometric shoulder flexion task against gravity, including arm weight. Seated participants had to hold a 2 kg dumbbell in front of them with the arm extended and pull up as much as possible for 3 s. The dumbbell was linked with a static rope to a force sensor attached to the ground. Participants had to stay in contact with the backrest during the entire contraction. Any help of the contralateral arm was prohibited. Experimenter encouraged the participant verbally. We retained the maximum score over 3 trials separated with a rest time of 1 min. Maximum antigravity shoulder torque was calculated based on the recorded maximum force and lever arm, plus the torque due to the weight of the arm during the task.

Experimental setup. Movements were recorded at 100 Hz with 8 infrared optical cameras from the Vicon Motion Capture System (Vantage V5, lens 8.5 mm, Oxford Metrics, UK). VICON Nexus 2 software was used to save time series of each marker. The experimenter placed markers on the target and the manubrium, and for each body side on the 1st metacarpal, the lateral epicondyle of the humerus, the acromion process, and the iliospinal. For each side, we corrected iliospinal marker position before data analysis to correspond to best to the anatomical centre of hips joints. Markers of both sides were needed to compute shoulder and trunk angles.

Kinematic analysis. Data analysis was performed with Scilab 6.0.2.

First, all position time series were low pass filtered at 5 Hz with a dual pass second order Butterworth filter.

Second, we computed the beginning and end of the reach. Because the goal of a reaching task is to bring the hand to the target, that is, to reduce the hand-to-target distance, what is important for task success is the hand-to-target Euclidean distance. The hand-to-target Euclidean distance summarises the 3D effector space into a 1D task space (where movement matters) leaving aside a 2D null space (where movement does not impact task success). We fixed the beginning of the movement (t_0) when the Euclidean velocity of the hand in task space became positive and remained positive until the maximum velocity. The end of the movement (t_{final}) was when the Euclidean distance to the target reached its minimum.

Third, we computed the trunk recruitment as a percentage of reach length. Reach length is the hand-to-target Euclidean distance at t_0 . In order to know the normalised implication of the trunk to the reach, we computed trunk recruitment as the ratio between shoulder displacement and reach length. More precisely, trunk recruitment = (shoulder to target distance at t_0 – shoulder to target distance at $t_{\text{final}})$ / (hand to target distance at t_0 – hand to target distance at t_{final}). Shoulder displacement is sensitive to both trunk anterior flexion and trunk rotation. Over the 5 trials, the trial with the median trunk recruitment was retained for data analysis.

Fourth, antigravity shoulder torque was computed as the Euclidean norm of the 3D static torque against gravity at the final posture. In order to compute the static torque, the positions of centre of mass and the absolute weight of upper limbs were approximated from the height and weight of participants following De Leva's equations (1996). Our model considers the torques applied to the shoulder by the upper arm (shoulder to elbow link), the forearm and hand (elbow to hand link) and the dumbbell, as described in the Eq. (1).




Figure 1. Schema of the four experimental conditions. In the loaded conditions (upper row), participants held a dumbbell which mass corresponded to 75% of their maximal voluntary shoulder force (MVF). In the unloaded conditions (lower row), participants had to reach the target without the dumbbell. In the spontaneous conditions (left column), participants had to reach the target in a natural way. In the trunk restraint conditions (right column), participants had to reach the target while minimising trunk movement.

$$\text{antigravity shoulder torque} = \sqrt{\bar{\tau}_{\text{upperarm}}^2 + \bar{\tau}_{\text{forearmHand}}^2 + \bar{\tau}_{\text{dumbbell}}^2} \quad (1)$$

The torque applied to the shoulder by each component of the model was calculated as the cross product of the lever arm vector and the gravity vector of the component (Eq. 2).

$$\bar{\tau}_{\text{component}} = \vec{r} \times \vec{F} \quad (2)$$

where \vec{r} is the lever arm vector (a position vector from the shoulder to the centre of mass of the component) and \vec{F} is the gravity vector of the component. The equation can be developed as:

$$\bar{\tau}_{\text{component}} = \left(\overrightarrow{\text{Position}}_{Cm} - \overrightarrow{\text{Position}}_{\text{shoulder}} \right) \times (\vec{F}_g * \text{Mass}) \quad (3)$$

where $\overrightarrow{\text{Position}}_{Cm}$ is the 3D position of the centre of mass of the component, $\overrightarrow{\text{Position}}_{\text{shoulder}}$ is the 3D position of the shoulder joint, \vec{F}_g is the force of gravity and *Mass* is the mass of the component.

Statistical analysis. The statistical analysis was conducted using R 3.6.1 with *rstatix* and *stats* packages. Density plots and quantiles/quantiles plots revealed a violation of the normality assumption inside data groups. Therefore, all hypotheses were tested in a non-parametric way, using paired samples Wilcoxon signed rank tests. Because of the multiple comparisons, *p*-values were corrected by the method of Benjamini and Hochberg (1995) with a false discovery rate (FDR) of 0.05. This method aims to control the expected proportion of false

discoveries amongst the rejected hypotheses (the FDR) in order to reduce type I errors while limiting type II errors. The FDR method is less stringent than the family-wise error rate and is therefore more powerful than a simple Bonferroni-type procedure. The significance level was set at 0.05 in all analyses. The effect size r was calculated as the Z statistic divided by the square root of the number of observations¹⁸. Data are presented as median (\pm interquartile range).

Results

Increasing the weight of the arm raises trunk recruitment. Reaching in the spontaneous unloaded condition induced a trunk recruitment of 11% of reach length. Trunk recruitment increased to 27.5% when loaded ($W=0$, $p<0.001$, $r=0.87$). In the trunk restraint conditions, trunk recruitment could be reduced to 8% when unloaded ($W=22$, $p<0.001$, $r=0.72$), and to 12% when loaded ($W=0$, $p<0.001$, $r=0.87$) (Fig. 2a). The increase in trunk recruitment was not different between subjects with the dominant side loaded and subjects with the non-dominant side loaded ($W=97$, $p=0.54$, $r=0.13$). These results indicate that handling a heavy dumbbell during reaching increases trunk recruitment (Table 1). When asked, all participants were able to reduce trunk recruitment in the loaded condition.

Trunk restraint reduces antigravity shoulder torque. Trunk restraint increased shoulder anti-gravity torque in the final posture from 11.7 to 12.7% of maximum when the hand is unloaded ($W=351$, $p<0.001$, $r=0.87$) and from 60.4 to 74.1% of maximum in the loaded condition ($W=349$, $p<0.001$, $r=0.86$) (Fig. 2b). The effect of trunk restraint on shoulder torque was not different between subjects with the dominant side loaded and subjects with the non-dominant side loaded ($W=92$, $p=0.72$, $r=0.08$). In other words, trunk restraint increased the mechanical cost that is needed at the shoulder to maintain the final posture. Given the increased arm weight in the loaded condition, we made no comparison between unloaded and loaded conditions.

Discussion

We tested the hypothesis that, when reaching to grasp an object from a seated position, people recruit the trunk when shoulder torque is too close to maximal. We found that when the hand is loaded to 75% of the maximum antigravity shoulder torque, trunk recruitment increases from 11.0 to 27.5%. By flexing and rotating the trunk, participants spontaneously reduced antigravity shoulder torque from 74.1 to 60.4% of maximum voluntary torque.

Trunk is a crucial component of the reaching coordination. The spontaneous unloaded condition informs about the final posture that is naturally selected when reaching to grasp an object from a seated position. From a side view, the task requires to move the hand forward. To do this, people combine elbow extension (43°) and shoulder flexion (42°), with no flexion of the trunk (0°). From a frontal view, the task requires moving the hand towards the midline of the body. To do this, people combine shoulder adduction (22°), with a small trunk rotation (4°) (see Table 1; Fig. 3, top row).

In the spontaneous unloaded condition, participants use a combination of elbow extension and shoulder flexion and adduction, but hardly recruit the trunk (Fig. 3, top row). This “arm strategy” is in agreement with the theory of optimal control, which says that human movement minimise costs³. Indeed, although participants could recruit the trunk, they do not because this would imply moving a large and heavy segment, and therefore to mobilise large muscles. It is more economical to mainly use the arm and forearm segments, which are less heavy and have less inertia than the trunk.

In the spontaneous loaded condition, participants still use a combination of elbow extension and shoulder flexion and adduction, but they do recruit the trunk (Fig. 3, bottom row). This “trunk strategy” decreases shoulder-elbow participation, and therefore minimises antigravity shoulder torque (Fig. 2). Indeed, when the arm is loaded and the trunk restricted, antigravity shoulder torque of the final posture is 74% of the maximum, which is a high intensity. According to Diedrichsen et al. (2010), when muscles are strongly activated, metabolic processes are not optimised and therefore the need for adenosine triphosphate (ATP) to maintain exertion is high. In addition, signal-dependent noise increases with the intensity of activation¹⁹, which may affect the precision of the reach. Thus, when the arm is loaded, the “arm strategy” likely involves a more energy-consuming and imprecise movement. Participants therefore change their coordination and their final posture so to reduce antigravity shoulder torque. The “trunk strategy” distributes the work over more effectors and thus reduces the load on shoulder muscles. The “trunk strategy” improves metabolic processes and reduces signal-dependent noise when the weight of the arm is high relative to the strength of the shoulder. This gain probably outweighs the metabolic expense required to recruit trunk muscles.

Diedrichsen et al. (2010) rename this principle “distribution of work across multiple effectors”. The authors consider that the cost of the task is equal to the squared sum of the motor commands. Indeed, this cost function is well correlated with ATP consumption and therefore with the effort required. According to this cost function, by penalising high activations, the distribution principle would minimise the total cost of the movement, and therefore minimise the required effort. Our results are consistent with the conclusions of Wang and Dounskaia (2012). They show that increasing arm weight accentuates, and thus makes visible the minimisation of muscular effort. Thus, when the arm is unloaded, effort minimisation is still at work but is less noticeable.

Here, we confirm the idea of Peeters et al. (2018) and Fuller et al. (2013) that trunk movement occurs when the shoulder has to develop a too great torque in relation to its capacity. This means that in addition to stabilising the arm and regulating the reachable distance^{8,21}, trunk movements emerge to avoid high shoulder torques.




Figure 2. Trunk recruitment (Panel A) and antigravity shoulder torque (Panel B) as a function of trunk restraint and hand load. Trunk recruitment is in percentage of reach length. Antigravity shoulder torque at the final posture is in percentage of maximal voluntary shoulder torque. In each boxplot, the internal horizontal bar represents the median, boxes extend from first to third quartile, whiskers extend from minimal to maximal values excluding outliers and white circles represent outliers. All comparisons are statistically significant. The figure shows that trunk recruitment stays low in the unloaded conditions. When the hand is loaded, trunk recruitment is doubled, and antigravity shoulder torque is decreased by 18% in the spontaneous condition.

Limitation. Participants did the reaching movement without holding a dumbbell in the unloaded condition. So, it is possible that some of the coordination differences are due to the difference in hand orientation between the dumbbell (loaded) and the non-dumbbell (unloaded) condition. However, in order to avoid this bias, the target was designed not to interfere with the movement performed with the dumbbell, and participants were instructed to hit the target with the side of their thumb, which theoretically produces a strictly similar movement, whether or not the participant is holding the dumbbell.

Trunk hand	Trunk restraint unloaded	Spontaneous unloaded	Trunk restraint loaded	Spontaneous loaded
Elbow extension (°)	43 (± 19)	37 (± 13)	29 (± 16)	14 (± 19)
Shoulder flexion (°)	42 (± 10)	41 (± 12)	39 (± 12)	31 (± 15)
Shoulder adduction (°)	22 (± 11)	19 (± 8)	11 (± 12)	12 (± 8)
Trunk flexion (°)	0 (± 0)	1 (± 0)	1 (± 2)	5 (± 5)
Trunk rotation (°)	4 (± 2)	6 (± 3)	12 (± 6)	17 (± 6)
Hand displacement (mm)	314 (± 46)	314 (± 42)	317 (± 51)	338 (± 62)
Shoulder displacement (mm)	25 (± 13)	36 (± 16)	63 (± 40)	138 (± 86)
Trunk displacement (mm)	5 (± 4)	11 (± 5)	38 (± 25)	79 (± 73)

Table 1. Differences in angular rotation and raw displacements between final and initial posture. Trunk displacement is the distance travelled by the manubrium: it captures mainly trunk anterior flexion. Shoulder displacement is the distance travelled by the acromion process: it is sensitive to both trunk anterior flexion and trunk rotation. When the hand is unloaded and the trunk is restrained, the coordination is mainly based on elbow and shoulder use (left of the table). When the hand is loaded or the trunk is free, trunk recruitment increases, which reduces elbow and shoulder use (right of the table).




Figure 3. Final posture of a typical participant in the spontaneous condition as a function of load (Unloaded: top row, Loaded: bottom row). Diagrams are viewed from the side (sagittal plane, left column), viewed from above (transverse plane, middle column) and viewed from the front (frontal plane, right column). The red circle represents the target. When the hand is loaded, the participant has less elbow extension, less shoulder flexion and shoulder abduction at the expense of greater trunk flexion and rotation. The figure shows that the greater recruitment of the trunk brings the shoulder closer to the target, resulting in less gravitational torque at the shoulder, as can be inferred by comparing the sagittal views.

Implications for motor control. The theory of optimal control states that movement coordination is performed to meet the constraints of the task while minimising a cost function²². Objective cost functions were first explored to minimise a mechanical cost^{23,24}, but according to Loeb (2012), cost functions not based on subjective feelings make no sense in a neurobiological system. More human-oriented functions were then developed, such as minimum neuromuscular effort²⁶, minimum metabolic cost^{27,28} or maximum comfort²⁹. Finally, other functions combine objective and subjective effort³⁰ or introduce modulation by individual traits^{4,31}. In the face of this diversity, there is currently no consensus on the list of ingredients to include in the cost function.

Here we show that in a seated reaching task, when the arm is heavy relative to the shoulder strength, healthy participants modify their final posture to avoid excessive antigravity torque at the shoulder. By recruiting the trunk in the loaded condition, the antigravity torque at the shoulder decreases from about 75–60% of maximum (Fig. 2, pink boxes). Although this result is consistent with the predictions of the optimal control framework, future work is necessary to ensure that the observed strategy is optimal.

Instead of performing optimally, biological organisms may just do “good enough”²⁵. Indeed, muscle coordination might be habitual rather than optimal in a wide variety of situations, which would explain the absence of consensus on the nature of the cost function. Following this approach, a disturbed sensorimotor system will more probably exhibit, at first, the habitual coordination adapted to the new constraints instead of the optimal solution that might come with learning³². In the present experiment, when loaded, participants immediately




Figure 4. Shoulder reserve in final posture as a function of trunk restraint and hand load. Shoulder reserve is in percentage of maximal antigravity shoulder torque. In each barplot, the top of the bar represents the median, error bar extends from the first to the third quartile. All comparisons are statistically significant. The figure shows that shoulder reserve stays high in the unloaded conditions. When the hand is loaded and the trunk is restrained, shoulder reserve is considerably reduced.

performed significant trunk recruitment. This suggests that participants have already learned that the habitual coordination is no longer “good enough” when the torque at the shoulder becomes excessive.

The strategy of coordination is directly adapted to the constraints to avoid a shoulder torque greater than about 60% of maximum (Fig. 2, panel B). If we consider the remaining shoulder torque as a shoulder torque reserve (i.e., $100 - 60\% = 40\%$), then it appears that participants reorganise their coordination to guarantee a shoulder torque reserve of about 40% (Fig. 4). In the trunk restraint condition, one would have expected that trunk recruitment would tend towards zero. Yet, participants chose to retain some trunk recruitment, resulting in 25% unmobilised shoulder reserve (Fig. 4, pink bars). This suggests that the spontaneous 40% reserve might be the sum of a mobilisable comfort reserve of 15%, and a non-mobilisable safety reserve of 25%. Keeping a reserve of torque is tantamount to the safety margin when grasping an object³³, and more generally, keeping a safety reserve is a behavioural strategy adopted by humans to be able to react to unexpected perturbations, which is mandatory to succeed in the task every time.

Although most of the current cost functions are linked to human effort, they are generally derived from experiments where participants spontaneously perform far from their limits^{31,34}. Here we demonstrate that, to better model the reaching movement where people are pushed towards their maximum, realistic cost functions should penalise too strong activations. Our results suggest that a reserve of about 40% of maximal force is the limit of acceptable activation without spontaneously recruiting a novel degree of freedom.

Limitations and future work. If this work lays the foundations for understanding the activation threshold beyond which the effort is avoided, it remains preliminary. First, the antigravity torque at the shoulder is only estimated statically for the final posture, which does not fully reflect the actual activation of the shoulder muscles. Torques could be estimated dynamically during the reach and muscle activations could be investigated using EMG analysis. However, with a motion time of 1.97 s (± 0.6) and a dwell time of at least 1 s at the target, the antigravitational torque at the final posture is the main constraint (e.g., an order of magnitude greater than the maximum dynamic torque³⁵). This results in greater activation of the anterior deltoid at the end of movement in slow reaches³⁶. Second, further experiments could measure how trunk recruitment increases as a function of shoulder torque, for example to get a picture of the non-linearity of this relation. Third, the reaching task studied here is very stereotypical. Further work could measure the evolution of coordination in more ecological tasks, with greater or lesser spatial accuracy requirements.

Perspectives for stroke research. Our result that trunk recruitment is a cost-effective solution when shoulder torque is too close to maximal has consequences for the interpretation of movement disorders. The most disabling factor after a stroke is muscle weakness in the hemiparetic arm³⁷. In other words, after a stroke, the weight of the arm is too heavy for the available strength of the shoulder muscles^{38,39}. As a result, people with stroke are close to maximum neuromuscular activation during a reaching movement⁴⁰. After a stroke, it is likely that the principles governing sensorimotor control and learning are the same^{41,42}. Hence, with the very same control policy in the face of an atypical body, the emerging coordination is atypical⁴³. By recruiting the trunk,

patients follow the typical control policy: they reduce the necessary shoulder torque and thus minimise effort. This interpretation is consistent with empirical results showing that arm weight support decreased neuromuscular activation of the shoulder^{44,45}, and so increased the reachable space of the patients^{46,47}. This confirms that the ability to lift the arm is a strong limiting factor when reaching after a stroke.

When the available force is limited, it may be more efficient to freely use all the available degrees of freedom instead of spending more attention, effort, energy to perform the task more “normally” (i.e. with the “arm strategy”). Thus, trunk compensation by stroke victims is not ineluctably a “bad synergy” resulting from a pathological synergy or from an impaired control^{48,49}, but could instead testify that stroke movement continue to follow optimal control principles, but with different constraints (such as reduced maximum shoulder torque or greater noise), as claimed in other diseases⁴³. Further research is needed to confirm that both people with and without stroke recruit their trunk when the arm is too heavy for the available shoulder force. For example, experimentally reducing the weight of the arm of post-stroke individuals should result in a proportional reduction in trunk compensation. If so, increasing antigravity shoulder force might be a way to facilitate arm use in individuals with stroke. Thus, it may be worth considering training that emphasises strength in addition to conventional therapies⁵⁰.

However, care must be taken not to confuse the strength reserve with what we call a “coordination reserve”. The latter is a form of neuroplastic cognitive reserve that explains “the disjunction between the degree of brain damage and its clinical outcome”⁵¹ and that can be drawn upon to change the coordination through learning^{52,53}.

Conclusion

Our experiment shows that, in young healthy persons reaching at a target while seated, trunk recruitment becomes visible when the participant faces antigravity force limitations at the shoulder. The recruitment of the trunk is a cost-effective solution to keep the activation of the shoulder muscles under 60% of their maximum. More generally, our results suggest that all joints in the kinematic chain actively participate in the movement, but that the amplitude of their recruitment is mediated by force constraints: energy-consuming joints are recruited only when necessary, to minimise the use of the force reserve of the other joints.

Data availability

The raw datasets generated during the current study are available in the Open Science Framework repository: <https://osf.io/r3xcu/>.

Code availability

The code generated during the current study is available in the Open Science Framework repository: <https://osf.io/r3xcu/>.

Received: 16 April 2021; Accepted: 12 November 2021

Published online: 26 November 2021

References

- Bernstein. *The Co-ordination and Regulation of Movements*. (Pergamon Press, 1967).
- Jeannerod, M. Intersegmental coordination during reaching at natural visual objects. *Attent. Perform.* **IX**, 153–169 (1981).
- Todorov, E. & Jordan, M. I. Optimal feedback control as a theory of motor coordination. *Nat. Neurosci.* **5**, 1226–1235 (2002).
- Rigoux, L. & Guigon, E. A model of reward- and effort-based optimal decision making and motor control. *PLOS Comput. Biol.* **8**, e1002716 (2012).
- Diedrichsen, J., Shadmehr, R. & Ivry, R. B. The coordination of movement: optimal feedback control and beyond. *Trends Cogn. Sci.* **14**, 31–39 (2010).
- Biomechanics of Anthropomorphic Systems*. vol. 124 (Springer, 2019).
- Prilutsky, B. I. & Zatsiorsky, V. M. Optimization-based models of muscle coordination. *Exerc. Sport Sci. Rev.* **30**, 32 (2002).
- Kaminski, T. R., Bock, C. & Gentile, A. M. The coordination between trunk and arm motion during pointing movements. *Exp. Brain Res.* **106**, 457–466 (1995).
- Cirstea, M. C. & Levin, M. F. Compensatory strategies for reaching in stroke. *Brain* **123**, 940–953 (2000).
- Levin, M. F., Liebermann, D. G., Parmet, Y. & Berman, S. Compensatory versus noncompensatory shoulder movements used for reaching in stroke. *Neurorehabil. Neural Repair* **30**, 635–646 (2016).
- Peeters, L. H. C., de Groot, I. J. M. & Geurts, A. C. H. Trunk involvement in performing upper extremity activities while seated in neurological patients with a flaccid trunk—A review. *Gait Posture* **62**, 46–55 (2018).
- Fuller, J. R., Fung, J. & Côté, J. N. Posture-movement responses to stance perturbations and upper limb fatigue during a repetitive pointing task. *Hum. Mov. Sci.* **32**, 618–632 (2013).
- Dysart, M. J. & Woldstad, J. C. Posture prediction for static sagittal-plane lifting. *J. Biomech.* **29**, 1393–1397 (1996).
- Bakhti, K. K. A., Mottet, D., Schweighofer, N., Froger, J. & Laffont, I. Proximal arm non-use when reaching after a stroke. *Neurosci. Lett.* **657**, 91–96 (2017).
- Bakhti, K. K. A., Laffont, I., Muthalib, M., Froger, J. & Mottet, D. Kinect-based assessment of proximal arm non-use after a stroke. *J. NeuroEng. Rehabil.* **15**, 104 (2018).
- De Leva, P. Adjustments to Zatsiorsky–Seluyanov’s segment inertia parameters. *J. Biomech.* **29**, 1223 (1996).
- Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. Roy. Stat. Soc.: Ser. B (Methodol.)* **57**, 289–300 (1995).
- R. Rosenthal, H. Cooper, & L. Hedges. Parametric measures of effect size. in *The handbook of research synthesis*. 231–244 (1994).
- Harris, C. M. & Wolpert, D. M. Signal-dependent noise determines motor planning. *Nature* **394**, 780 (1998).
- Wang, W. & Dounskaia, N. Load emphasizes muscle effort minimization during selection of arm movement direction. *J. Neuroeng. Rehabil.* **9**, 70 (2012).
- Robertson, J. V. G. & Roby-Brami, A. The trunk as a part of the kinematic chain for reaching movements in healthy subjects and hemiparetic patients. *Brain Res.* **1382**, 137–146 (2011).
- Todorov, E. Optimality principles in sensorimotor control. *Nat. Neurosci.* **7**, 907 (2004).
- Nelson, W. L. Physical principles for economies of skilled movements. *Biol. Cybern.* **46**, 135–147 (1983).

24. Uno, Y., Kawato, M. & Suzuki, R. Formation and control of optimal trajectory in human multijoint arm movement. *Biol. Cybern.* **61**, 89–101 (1989).
25. Loeb, G. E. Optimal isn't good enough. *Biol. Cybern.* **106**, 757–765 (2012).
26. Guigon, E., Baraduc, P. & Desmurget, M. Computational motor control: Redundancy and invariance. *J. Neurophysiol.* **97**, 331–347 (2007).
27. Praagman, M., Chadwick, E. K. J., van der Helm, F. C. T. & Veeger, H. E. J. The relationship between two different mechanical cost functions and muscle oxygen consumption. *J. Biomech.* **39**, 758–765 (2006).
28. Shadmehr, R., Huang, H. J. & Ahmed, A. A. A representation of effort in decision-making and motor control. *Curr. Biol.* **26**, 1929–1934 (2016).
29. Cruse, H., Wischmeyer, E., Brüwer, M., Brockfeld, P. & Dress, A. On the cost functions for the control of the human arm movement. *Biol. Cybern.* **62**, 519–528 (1990).
30. Wang, C., Xiao, Y., Burdet, E., Gordon, J. & Schweighofer, N. The duration of reaching movement is longer than predicted by minimum variance. *J. Neurophysiol.* **116**, 2342–2345 (2016).
31. Berret, B., Castanier, C., Bastide, S. & Deroche, T. Vigour of self-paced reaching movement: cost of time and individual traits. *Sci. Rep.* **8**, 1–14 (2018).
32. de Rugy, A., Loeb, G. E. & Carroll, T. J. Muscle coordination is habitual rather than optimal. *J. Neurosci.* **32**, 7384–7391 (2012).
33. Danion, F. Grip force safety margin in rock climbers. *Int. J. Sports Med.* **29**, 168–172 (2008).
34. Berret, B., Chiovetto, E., Nori, F. & Pozzo, T. Evidence for composite cost functions in arm movement planning: An inverse optimal control approach. *PLOS Comput. Biol.* **7**, e1002183 (2011).
35. Olesh, E. V., Pollard, B. S. & Gritsenko, V. Gravitational and dynamic components of muscle torque underlie tonic and phasic muscle activity during goal-directed reaching. *Front. Hum. Neurosci.* **11**, 474 (2017).
36. Papaxanthis, C., Pozzo, T. & Schieppati, M. Trajectories of arm pointing movements on the sagittal plane vary with both direction and speed. *Exp. Brain Res.* **148**, 498–503 (2003).
37. Canning, C. G., Ada, L., Adams, R. & O'Dwyer, N. J. Loss of strength contributes more to physical disability after stroke than loss of dexterity. *Clin. Rehabil.* **18**, 300–308 (2004).
38. Ada, D. L., O'Dwyer, N., Ada, D. L., O'Dwyer, N. & O'Neill, E. Relation between spasticity, weakness and contracture of the elbow flexors and upper limb activity after stroke: An observational study. *Disabil. Rehabil.* **28**, 891–897 (2006).
39. Bourbonnais, D., Vanden Noven, S. Weakness in patients with hemiparesis. *Am. J. Occup. Ther.* **43**, 313–319 (1989).
40. McCrea, P. H., Eng, J. J. & Hodgson, A. J. Saturated muscle activation contributes to compensatory reaching strategies after stroke. *J. Neurophysiol.* **94**, 2999–3008 (2005).
41. Kitago, T. & Krakauer, J. W. Motor learning principles for neurorehabilitation. in *Handbook of clinical neurology*. vol. 110, 93–103 (Elsevier, 2013).
42. Mottet, D., van Dokkum, L. E. H., Froger, J., Gouaich, A. & Laffont, I. Trajectory formation principles are the same after mild or moderate stroke. *PLoS ONE* **12**, e0173674 (2017).
43. Latash, M. L. & Anson, J. G. What are “normal movements” in atypical populations?. *Behav. Brain Sci.* **19**, 55–68 (1996).
44. Coscia, M. *et al.* The effect of arm weight support on upper limb muscle synergies during reaching movements. *J. NeuroEng. Rehabil.* **11**, 22 (2014).
45. Runnalls, K. D., Ortega-Auriol, P., McMorland, A. J. C., Anson, G. & Byblow, W. D. Effects of arm weight support on neuromuscular activation during reaching in chronic stroke patients. *Exp. Brain Res.* **237**, 3391–3408 (2019).
46. Dewald, J. P. A., Sheshadri, V., Dawson, M. L. & Beer, R. F. Upper-limb discoordination in hemiparetic stroke: Implications for neurorehabilitation. *Top. Stroke Rehabil.* **8**, 1–12 (2001).
47. Sukal, T. M., Ellis, M. D. & Dewald, J. P. A. Shoulder abduction-induced reductions in reaching work area following hemiparetic stroke: Neuroscientific implications. *Exp. Brain Res.* **183**, 215–223 (2007).
48. Dewald, J. P. A., Pope, P. S., Given, J. D., Buchanan, T. S. & Rymer, W. Z. Abnormal muscle coactivation patterns during isometric torque generation at the elbow and shoulder in hemiparetic subjects. *Brain* **118**, 495–510 (1995).
49. Subramanian, S. K. *et al.* Motor-equivalent intersegmental coordination is impaired in chronic stroke. *Neurorehabil. Neural Repair* **34**, 210–221 (2020).
50. Hill, T. R. *et al.* Maximal strength training enhances strength and functional performance in chronic stroke survivors. *Am. J. Phys. Med. Rehabil.* **91**, 393–400 (2012).
51. Stern, Y. Cognitive reserve. *Neuropsychologia* **47**, 2015–2028 (2009).
52. Cramer, S. C. *et al.* Harnessing neuroplasticity for clinical applications. *Brain* **134**, 1591–1609 (2011).
53. Winstein, C. J. & Kay, D. B. Translating the science into practice. in *Progress in brain research* vol. 218, 331–360 (Elsevier, 2015).

Acknowledgements

This study was supported by the LabEx NUMEV (ANR-10-LABX-0020) within the I-SITE MUSE. We thank Victor Barradas and Nicolas Schweighofer for their comments on the article. We acknowledge Germain Faily for drawing the schemata of the experimental conditions in Figs. 1 and 3.

Author contributions

G.F. designed the protocol, recorded and analysed the data, wrote the first version of the manuscript. D.M. and J.F. assisted G.F. in designing the protocol, analysing the data and provided guidance for improving the manuscript. S.P. assisted G.F. in setting up the protocol, recording the data and improving the manuscript. All authors accepted the latest version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to D.M.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021